

# ON BURGHELEA CONJECTURE

ALEXANDER DRANISHNIKOV

**ABSTRACT.** We prove the Burghelea Conjecture for all groups with finite cohomological dimension satisfying some additional condition on the string cohomology.

## 1. INTRODUCTION

We recall that for an algebra  $A$  over a field  $k$  (generally, over a commutative ring)  $HH_*(A)$  and  $HC_*(A)$  denote the Hochschild homology and the cyclic homology of  $A$  [L],[R]. These two homology groups are connected by the Connes exact sequence

$$\cdots \rightarrow HC_{n-1}(A) \rightarrow HH_n(A) \rightarrow HC_n(A) \xrightarrow{S} HC_{n-2}(A) \rightarrow \cdots$$

We use notation  $PHC_*(A)$  for the periodic cyclic homology,  $*$  = 0, 1. The shift homomorphism  $S$  in the Connes exact sequence defines the inverse sequence  $\{HC_{*+2n}, S\}_{n \in \mathbb{N}}$ . We note that for the periodic cyclic homology there is an epimorphism  $PHC_*(A) \rightarrow \lim_{\leftarrow} \{HC_{*+2n}(A), S\}$  with the  $\lim^1$ -kernel. The kernel is trivial if the Mittag-Leffler condition is satisfied in particular when  $HH_m(A) = 0$  for all sufficiently large  $m$  (see [R] Corollary 6.1.23).

We consider the case when  $k = \mathbb{Q}$  and  $A$  is the group algebra  $\mathbb{Q}G$  for a discrete group  $G$ . It was known [ML] well before the invention of the cyclic homology in early 80s ([C],[LQ],[T]) that  $HH_n(\mathbb{Q}G) = H_n(G, \widetilde{\mathbb{Q}G})$  where  $\widetilde{\mathbb{Q}G}$  is the  $G$ -module  $\mathbb{Q}G$  defined by the  $G$ -action on  $G$  by conjugation. Let  $\langle G \rangle$  denote the set of conjugacy classes on  $G$  and let  $G_x$  denote the centralizer of  $x \in [x] \in \langle G \rangle$ . Clearly,  $G_y \cong G_z$  for  $y, z \in [x]$ . Since

$$\widetilde{\mathbb{Q}G} = \bigoplus_{[x] \in \langle G \rangle} \mathbb{Q}(G/G_x) = \bigoplus_{[x] \in \langle G \rangle} \mathbb{Q} \otimes_{G_x} \mathbb{Q}G,$$

---

*Date:* February 21, 2017.

2000 *Mathematics Subject Classification.* Primary 20G10; Secondary 16E40, 55P50 .

The author was supported by NSF DMS-1304627.

we obtain

$$HH_n(\mathbb{Q}G) = \bigoplus_{[x] \in \langle G \rangle} H_n(G, \mathbb{Q} \otimes_{G_x} \mathbb{Q}G).$$

By the Shapiro Lemma [Br],  $H_n(G, \mathbb{Q} \otimes_{G_x} \mathbb{Q}G) = H_n(G_x, \mathbb{Q})$ . Thus,

$$HH_n(\mathbb{Q}G) = \bigoplus_{[x] \in \langle G \rangle} H_n(G_x, \mathbb{Q}).$$

Burghelea have computed cyclic homology of group algebras [Bu]

$$HC_n(\mathbb{Q}G) = \bigoplus_{\langle G \rangle^{fin}} [H_*(N_x, \mathbb{Q}) \otimes H_*(\mathbb{C}P^\infty; \mathbb{Q})]_n \oplus \bigoplus_{\langle G \rangle^\infty} H_n(N_x, \mathbb{Q})$$

where  $N_x = G_x / \langle x \rangle$  is the reduced centralizer,  $\langle G \rangle^{fin}$  is the subset of  $\langle G \rangle$  of conjugacy classes of elements of finite order, and  $\langle G \rangle^\infty$  is the subset of conjugacy classes of elements of infinite order. This computation brought the following formula for the periodic cyclic homology of  $\mathbb{Q}G$  [Bu]

$$PHC_0(\mathbb{Q}G) = \bigoplus_{[x] \in \langle G \rangle^{fin}, n \geq 0} H_{2n}(N_x, \mathbb{Q}) \oplus \bigoplus_{\langle G \rangle^\infty} T_0(x, \mathbb{Q})$$

$$PHC_1(\mathbb{Q}G) = \bigoplus_{[x] \in \langle G \rangle^{fin}, n \geq 0} H_{2n+1}(N_x, \mathbb{Q}) \oplus \bigoplus_{\langle G \rangle^\infty} T_1(x, \mathbb{Q}).$$

Here the group  $T_*(x, \mathbb{Q})$  fits in the following short exact sequence

$$0 \rightarrow \varprojlim^1 \{H_{*-1+2n}(N_x, \mathbb{Q})\} \rightarrow T_*(x, \mathbb{Q}) \rightarrow \varprojlim \{H_{*+2n}(N_x, \mathbb{Q})\} \rightarrow 0$$

where bonding maps in the inverse sequences are the Gysin homomorphisms  $S : H_m(N_x, \mathbb{Q}) \rightarrow H_{m-2}(N_x, \mathbb{Q})$  coming from the fibration  $S^1 \rightarrow BG_x \rightarrow BN_x$ .

In the end of his paper [Bu] Burghelea stated the following conjecture:

**1.1. Conjecture (Burghelea).** *If  $K(G, 1)$  has the homotopy type of a finite CW-complex, then  $T_*(x, \mathbb{Q}) = 0$  for all  $x \in \langle G \rangle^\infty$ .*

Burghelea proved his conjecture for the fundamental groups of Riemannian manifolds with nonpositive sectional curvature. He also noticed that not every group satisfies the condition  $T_*(x, \mathbb{Q}) = 0$  for all  $x \in \langle G \rangle^\infty$ . We say that a group  $G$  satisfies the Burghelea Conjecture if it satisfies this condition for all  $x \in \langle G \rangle^\infty$ . Eckmann proved [E] the Burghelea Conjecture for groups  $G$  with  $hd_{\mathbb{Q}}G < \infty$  in the case when  $G$  is nilpotent or torsion free solvable group. Also he proved the Burghelea Conjecture for linear groups over fields of characteristic 0 and for groups with  $cd_{\mathbb{Q}}G \leq 2$ . R. Ji [J] proved the Burghelea Conjecture

for hyperbolic groups, arithmetic groups, and groups of polynomial growth.

In this paper we introduce some property of groups called property E (see Definition 2.3) and prove the following theorem.

**1.2. Theorem.** *Suppose that a group  $G$  satisfies  $cd(G) < \infty$  and all reduced centralizers  $N_x$ ,  $x \in G$ , have property E. Then  $T_*(x, \mathbb{Q}) = 0$  for all  $x \in G$ .*

Eckmann [E] (see also [Em]) noticed that the Hattori-Stallings rank map

$$r : K_0(\mathbb{Q}G) \rightarrow HC_0(\mathbb{Q}G) = \mathbb{Q}G/[\mathbb{Q}G, \mathbb{Q}G] = \bigoplus_{\langle G \rangle} \mathbb{Q}$$

(see [Br] for the definition) can be factored through the Connes-Karubi character

$$ch^n : K_0(kG) \rightarrow HC_{2n}(kG), \quad n \geq 0,$$

as  $r = S^n \circ ch^n$  where  $S^n = S \circ \dots \circ S$  is the  $n$  times iterated shift homomorphism. It follows from a result of Linnel [Li], that the Hattori-Stallings rank takes value in  $\mathbb{Q} \oplus \bigoplus_{\langle G \rangle^\infty} \mathbb{Q}$  where the first summand corresponds to the conjugacy class of the unit  $e \in G$ . Therefore, the Hattori-Stallings rank factors through  $\mathbb{Q} \oplus \bigoplus_{\langle G \rangle^\infty} T_0(x, \mathbb{Q})$ . Thus, for fixed  $G$  the Burghelea Conjecture implies the Bass Conjecture [Ba] on vanishing the terms of the image of the Hattori-Stallings map in  $\bigoplus_{\langle G \rangle^\infty} \mathbb{Q}$ .

Note that in its turn the Bass Conjecture for  $G$  implies the Idempotents Conjecture [Em]: *In the group algebra  $\mathbb{Q}G$  for a torsion free group  $G$  the equation  $x^2 = x$  has only trivial solutions  $x = 0$  and  $x = 1$ .*

The author is thankful to Alex Engel, Michal Marcinkowski, and Oscar Randal-Williams for pointing out on mistakes in the early versions of the paper and giving valuable remarks.

## 2. FREE LOOP SPACES

Let  $LX$  denote the free loop space on  $X$ ,  $LX = Map(S^1, X)$ . By  $L_0X$  we denote the space of free null-homotopic loops on  $X$ ,  $L_0X \subset LX$ . When  $X$  is simply connected we have  $L_0X = LX$ . The group  $S^1$  naturally acts on  $LX$  by  $(zf)(u) = f(z^{-1}u)$ . Let  $S(X)$  denote the orbit space  $LX/S^1$ , the space of 'strings'. Note that  $L_0X$  is  $S^1$ -invariant and denote by  $S_0(X)$  the orbit space  $L_0X/S^1$ , the space of null-homotopic 'strings' on  $X$ .

There is the evaluation fibration  $ev : LX \rightarrow X$ ,  $ev(\phi : S^1 \rightarrow X) = \phi(1)$ . Note that  $ev$  admits a section  $s : X \rightarrow LX$  and the fiber of  $ev$  is the loop space  $\Omega X$ .

We note that  $L$ ,  $L_0$ ,  $S$ , and  $S_0$  are covariant functors on the category of topological spaces. Each of these functors  $F$  takes a homotopy to a homotopy in a sense that there is a natural embedding  $F(X) \times I \rightarrow F(X \times I)$ .

Suppose that  $X$  is a proper metric space which uniformly locally path connected, i.e. there are  $\epsilon > 0$  and a continuous map  $\Phi : W_\epsilon \times [0, 1] \rightarrow X$  of the  $\epsilon$ -neighborhood  $W_\epsilon$  of the diagonal  $\Delta(X) \subset X \times X$  such that  $\Phi(x, y, -) : [0, 1] \rightarrow X$  is a path from  $x$  to  $y$  and  $\Phi(x, x, -)$  is a constant path for all  $(x, y) \in W_\epsilon$ . Then  $L_0X$  is uniformly locally path connected by means of the function  $\tilde{\Phi} : \tilde{W}_\epsilon \times [0, 1] \rightarrow L_0X$  defined as  $\tilde{\Phi}(f_1, f_2, t)(u) = \Phi(f_1(u), f_2(u), t)$  where  $\tilde{W}_\epsilon$  is the epsilon neighborhood of the diagonal  $\Delta(L_0X)$  in the uniform metric. We note that  $\tilde{W}_\epsilon$  is invariant with respect to the diagonal  $S^1$ -action and the map  $\tilde{\Phi}$  is equivariant. Then the space  $S_0(X)$  is locally contractible. Moreover, the space  $S_0(X)/X$  is locally contractible. Note that for such spaces the singular cohomology behaves well and agrees with the Čech cohomology [Bre].

We note that  $X$  is embedded in  $LX$  as well in  $S_0X$  via the constant loops. Let  $j_X : X \rightarrow S_0(X)$  denote that embedding. Note that  $j_X : 1 \rightarrow S_0$  is a natural transformation of functors.

**2.1. Proposition.** *For any space  $X$*

$$SX/X = (LX/X)/S^1 \quad \text{and} \quad S_0X/X = (L_0X/X)/S^1$$

*where we identify  $X$  with  $j_X(X)$ .*

*Proof.* This is an obvious statement on the set-theoretical level: Both spaces consists of orbits of non-constant loops  $f : S^1 \rightarrow X$  plus the special point  $* = \{X\}$ . The rest of the proof is an exercise on the quotient topology which we feel obligated to do since the spaces involved are not compact.

We recall that the main theorem on quotient map which states that for a quotient map  $p : X \rightarrow Y$  and a continuous map  $g : X \rightarrow Z$  constant on each point preimage  $p^{-1}(y)$  there is a continuous map  $f : Y \rightarrow Z$  such that  $f \circ p = g$ . We apply it to the quotient map  $p : LX/X \rightarrow (LX/X)/S^1$  and  $g : LX/X \rightarrow SX/X$  and to the quotient map  $LX \rightarrow LX/X$  and a map  $LX \rightarrow LX/X \rightarrow (LX/X)/S^1$  to obtain a map  $f : (LX/X)/S^1 \rightarrow SX/X$  and its inverse  $f^{-1} : SX/X \rightarrow (LX/X)/S^1$ . Here we used the fact that the composition of quotient maps is a quotient map.  $\square$

We consider the Borel construction [Bo] for the  $S^1$  action on  $L_0X/X$  and on  $S^{2N+1}$  for sufficiently large  $N$ :

$$\begin{array}{ccccc} S^{2N+1} & \longleftarrow & S^{2N+1} \times (L_0X/X) & \longrightarrow & L_0X/X \\ \downarrow & & \downarrow & & \downarrow \\ \mathbb{C}P^N & \xleftarrow{p_1} & S^{2N+1} \times_{S^1} (L_0X/X) & \xrightarrow{p_2} & S_0(X)/X. \end{array}$$

Every orbit  $S^1 z \in S_0(X)/X$  is homeomorphic to  $S^1/G_z$  where  $G_z \subset S^1$  is the stabilizer of  $z$ , a closed subgroup of  $S^1$ . Then the fiber  $p_2^{-1}(z)$  is homeomorphic to  $S^{2N+1}/G_z$ . We note that the special fiber  $F = p_2^{-1}(\{X\})$  of  $p_2$  is homeomorphic to  $\mathbb{C}P^N$ . We denote by

$$E(N, X) = (S^{2N+1} \times_{S^1} (L_0X/X))/F$$

the quotient space and by  $\bar{p}_2 : E(N, X) \rightarrow S_0(X)/X$  the induced map. Thus  $p_2 = \bar{p}_2 \circ q$  where  $q$  is the map collapsing  $F$ .

**2.2. Proposition.** *For a uniformly locally path connected proper metric space  $X$  the projection  $\bar{p}_2$  induces an isomorphism*

$$(\bar{p}_2)_* : H_i(E(N, X); \mathbb{Q}) \rightarrow H_i(S_0(X)/X; \mathbb{Q})$$

*in dimensions  $i \leq 2N$ .*

*Proof.* Since the spaces here are locally nice, it suffices to prove the statement for the cohomology. We consider the Leray spectral sequence of  $\bar{p}_2$ . Since the map  $\bar{p}_2$  is proper, the sequence converges to  $H^*(E(N, X); \mathbb{Q})$  [Bre]. Note that  $\bar{p}_2$  has one exceptional fiber which is a point. All other fibers are lens spaces  $L_m^{2N+1}$  possibly degenerated to the sphere  $S^{2N+1}$ . Therefore, in the Leray spectral sequence we have trivial stalks

$$\mathcal{H}^q(\bar{p}_2) = \lim_{\rightarrow} H^q(\bar{p}_2^{-1}(U); \mathbb{Q}) = H^q(\bar{p}_2^{-1}(x); \mathbb{Q}) = 0$$

for  $0 < q \leq 2N$ . Hence,  $E_2^{p,q} = 0$  for  $0 < q \leq 2N$  and the result follows.  $\square$

**2.3. Definition.** We say that a group  $\pi$  *satisfies property E* if the map  $E\pi/E\pi^{(n)} \rightarrow B\pi/B\pi^{(n)}$  defined by the universal covering  $p : E\pi \rightarrow B\pi$  induces a rational epimorphism of homotopy groups  $\pi_i(E\pi/E\pi^{(n)}) \otimes \mathbb{Q} \rightarrow \pi_i(B\pi/B\pi^{(n)}) \otimes \mathbb{Q}$  for  $i < 2n - 1$ .

**2.4. Proposition.** *Let  $X = B\pi/B\pi^{(n-1)}$ , for a group that satisfies property E. Then*

$$H_i(LX/X; \mathbb{Q}) = 0$$

*for  $0 < i < n - 1$  and  $n - 1 < i < 2n - 2$ .*

*Proof.* Consider the evaluation fibration  $ev : LX \rightarrow X$ . We claim that the rationalization of the fiber  $\Omega X$  has the rational homology groups of a wedge of  $(n-1)$ -spheres.

It is easy to show that

$$\pi_i(E\pi/E\pi^{(n-1)}) \otimes \mathbb{Q} = 0$$

for  $n < i < 2n-1$ . The property E implies that

$$\pi_i(B\pi/B\pi^{(n-1)}) \otimes \mathbb{Q} = 0$$

for  $n < i < 2n-1$ .

Clearly,  $\pi_i(B\pi/B\pi^{(n-1)}) = 0$  for  $i < n$ . Then the rationalization  $X_0$  of  $X$  has trivial homotopy groups in dimensions  $i < n$  and  $n < i < 2n-1$ . Then the rationalization of the loop space  $\Omega(X)_0 = \Omega(X_0)$  has trivial homotopy groups in dimensions  $i < n-1$  and  $n-1 < i < 2n-2$ .

Thus in the homology Leray-Serre spectral sequence of  $ev$  we have  $E_{p,q}^2 = 0$  for  $0 < q < n-1$  and  $n-1 < q < 2n-2$ . Since  $X$  is  $(n-1)$ -connected,  $E_{p,0}^2 = 0$  for  $0 < p < n$ . This implies that  $H_i(LX; \mathbb{Q}) = 0$  for  $0 < i < n-1$  and  $n < i < 2n-2$ . The only nonzero entry in the antidiagonal  $E_{p,q}^2$  with  $p+q = n$  is  $H_n(X; \mathbb{Q})$ . Since  $ev$  has a section, this entry lives to  $E^\infty$ . Hence, the inclusion  $X \rightarrow LX$  induces an isomorphism  $H_n(X; \mathbb{Q}) = H_n(LX; \mathbb{Q})$ . The splitting

$$H_i(LX) = H_i(X) \oplus H_i(LX/X)$$

generated by the section  $s$  implies that  $H_i(LX/X; \mathbb{Q}) = 0$  for  $0 < i < n-1$  and  $n-1 < i < 2n-2$ .  $\square$

**2.5. Proposition.** *Let  $\phi_{*,*} : E_{*,*}(p') \rightarrow E_{*,*}(p)$  be the morphism of the Leray-Serre spectral sequences generated by a morphism  $(f', f) : p' \rightarrow p$  of fiber bundles,  $p' : X' \rightarrow Y'$  and  $p : X \rightarrow Y$ .*

(a) *If  $\phi_{0,n}^2 : E_{0,n}^2(p') \rightarrow E_{0,n}^2(p)$  is surjective then also is  $\phi_{0,n}^\infty : E_{0,n}^\infty(p') \rightarrow E_{0,n}^\infty(p)$ .*

(b) *If all  $\phi_{k,l}^\infty : E_{k,l}^\infty(p') \rightarrow E_{k,l}^\infty(p)$  for  $k+l = n$  are surjective, then  $f'_* : H_n(X') \rightarrow H_n(X)$  is surjective.*

*Proof.* (a) We show by induction on  $r$  that  $\phi_{0,n}^r$  is surjective. Indeed, since in the commutative diagram

$$\begin{array}{ccccccc} \xrightarrow{d^r} & E_{0,n}^r(p') & \longrightarrow & E_{0,n}^{r+1}(p') & \longrightarrow & 0 \\ & \phi_{0,n}^r \downarrow & & \phi_{0,n}^{r+1} \downarrow & & \\ \xrightarrow{d^r} & E_{0,n}^r(p) & \xrightarrow{\psi} & E_{0,n}^{r+1}(p) & \longrightarrow & 0 \end{array}$$

$\phi_{0,n}^r$  and  $\psi$  are surjective, so is  $\phi_{0,n}^{r+1}$ .

(b) The homology groups  $H_n(X)$  is obtained from  $E_{n,0}^\infty(p)$  by consecutive extensions by  $E_{n-1,1}^\infty(p)$  then by  $E_{n-2,2}^\infty(p)$  and so on up to  $E_{0,n}^\infty$ . The same holds true for  $H_n(X')$ . Let  $A_0 = E_{n,0}^\infty(p)$ ,  $A'_0 = E_{n,0}^\infty(p')$  and let  $A_i$  and  $A'_i$  denote the corresponding intermediate extensions,  $i = 1, \dots, n$ . Thus,  $H_n(X) = A_n$  and  $H_n(X') = A'_n$ . We apply the epimorphism version of Five Lemma recursively to the diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & E_{n-i,i}^\infty(p') & \longrightarrow & A'_i & \longrightarrow & A'_{i-1} \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & E_{n-i,i}^\infty(p) & \longrightarrow & A_i & \longrightarrow & A_{i-1} \longrightarrow 0 \end{array}$$

to obtain the result.  $\square$

The main result of this section, Theorem 2.6, is rather technical. It is a computation of the string homology in some special case. Since we are doing it by rather elementary means we don't involve here the loop homology and the string homology theories [CJY],[We].

**2.6. Theorem.** *Suppose that a group  $\pi$  has property E and the classifying space  $B\pi$  of  $\pi$  is locally finite. Then the inclusion*

(a)  $j_\pi : B\pi \rightarrow S_0(B\pi)$  *induces an isomorphism*

$$(j_\pi)_* : H_n(B\pi; \mathbb{Q}) \rightarrow H_n(S_0(B\pi); \mathbb{Q})$$

*of the rational homology groups and*

(b) *for  $n \geq 4$  the inclusion homomorphism*

$$H_n(S_0(B\pi^{(n-2)}); \mathbb{Q}) \rightarrow H_n(S_0(B\pi); \mathbb{Q})$$

*is zero.*

*Proof.* (a) We show that  $H_*(S_0(B\pi), j_\pi(B\pi); \mathbb{Q}) = 0$ .

Note that the fiber  $\Omega(B\pi)$  of the evaluation fibration  $ev : L(B\pi) \rightarrow B\pi$  is homotopy equivalent to  $\pi$ . This fibration admits a section  $s$  and the path component of  $s(B\pi)$  is exactly  $L_0(B\pi)$ . Thus, the restriction  $p_0$  of  $ev$  to  $L_0(B\pi)$  is a fibration with homotopy trivial fiber. Hence the inclusion  $B\pi \rightarrow L_0(B\pi)$  is a homotopy equivalence. Thus,  $L_0(B\pi)/B\pi$  is contractible. Therefore the projection in the Borel construction

$$p_1 : S^{2N+1} \times_{S^1} (L_0(B\pi)/B\pi) \rightarrow \mathbb{C}P^N$$

induces isomorphism of homology groups. By Proposition 2.2 the projection

$$\bar{p}_2 : (S^{2N+1} \times_{S^1} (L_0(B\pi)/B\pi))/F \rightarrow S_0(B\pi)/j_\pi(B\pi)$$

induces isomorphisms of rational homology in dimensions  $\leq 2N$ . The exceptional fiber  $F$  of  $p_2$  is homeomorphic to  $\mathbb{C}P^N$ . It defines a section

of  $p_1$ . Thus collapsing  $F$  to a point defines a contractible space. Therefore,  $H_i(S_0(B\pi), j_\pi(B\pi); \mathbb{Q}) = 0$  for  $i \leq 2N$ . Since  $N$  is arbitrary, the result follows.

(b) In view of the exact sequence of the pair  $(S_0(B\pi), S_0(B\pi^{(n-2)}))$  it suffices to show that the homomorphism

$$q_* : H_n(S_0(B\pi); \mathbb{Q}) \rightarrow H_n(S_0(B\pi)/S_0(B\pi^{(n-2)}); \mathbb{Q})$$

generated by the collapsing map

$$q : S_0(B\pi) \rightarrow S_0(B\pi)/S_0(B\pi^{(n-2)})$$

is injective. We note that the collapsing map  $\psi : B\pi \rightarrow B\pi/B\pi^{(n-2)}$  induces an equivariant map  $L_0(\psi) : L_0(B\pi) \rightarrow L_0(B\pi/B\pi^{(n-2)})$  and, hence, it defines a map  $s(\psi) : S_0(B\pi) \rightarrow S_0(B\pi/B\pi^{(n-2)})$ . Note that the map  $s(\psi)$  factors through  $q$ ,  $s(\psi) = \psi' \circ q$ . Thus, it suffices to show that  $s(\psi)$  is injective for  $n$ -homology. Since  $(j_\pi)_*$  and  $i$  are isomorphisms in the commutative diagram

$$\begin{array}{ccc} H_n(B\pi/B\pi^{(n-2)}; \mathbb{Q}) & \xrightarrow{j'_*} & H_n(S_0(B\pi/B\pi^{(n-2)}); \mathbb{Q}) \\ i \uparrow & & s(\psi) \uparrow \\ H_n(B\pi; \mathbb{Q}) & \xrightarrow{(j_\pi)_*} & H_n(S_0(B\pi); \mathbb{Q}), \end{array}$$

it suffices to show that  $j'_*$  is a monomorphism.

Let  $X = B\pi/B\pi^{(n-2)}$ . We need to show that

$$j'_* : H_n(X; \mathbb{Q}) \rightarrow H_n(S_0(X); \mathbb{Q})$$

is a monomorphism. The exact sequence of the pair  $(S_0(X), X)$  reduces the problem to showing that the collapsing map  $q : S_0(X) \rightarrow S_0(X)/X$  induces an epimorphism  $q_* : H_{n+1}(S_0(X); \mathbb{Q}) \rightarrow H_{n+1}(S_0(X)/X; \mathbb{Q})$ .

We consider the commutative diagram generated by Borel's constructions for  $S^1$ -action on  $LX$ ,  $LX/X$ , and  $S^{2N+1}$  with  $N > n$ .

$$\begin{array}{ccc} H_{n+1}(S^{2N+1} \times_{S^1} LX; \mathbb{Q}) & \xrightarrow{(p'_2)_*} & H_{n+1}(S_0 X; \mathbb{Q}) \\ \bar{q}_* \downarrow & & q_* \downarrow \\ H_{n+1}(S^{2N+1} \times_{S^1} (LX/X); \mathbb{Q}) & \xrightarrow{(p_2)_*} & H_{n+1}((S_0 X)/X; \mathbb{Q}). \end{array}$$

It suffices to show that both  $(p_2)_*$  and  $\bar{q}_*$  are surjective. By Proposition 2.2 the reduced projection

$$\bar{p}_2 : (S^{2N+1} \times_{S^1} (LX/X))/p_2^{-1}(*) \rightarrow S_0 X/X$$

induces isomorphism of the rational homology in dimensions  $\leq 2N$ . Here  $*$  =  $\{X\}$  is the orbit of the fixed point. The projection

$$p_1 : S^{2N+1} \times_{S^1} (LX/X) \rightarrow \mathbb{C}P^N$$

is a locally trivial bundle with the fiber  $LX/X$ . The fixed point  $*$  of the  $S^1$ -action on  $LX/X$  defines a section  $s$  of  $p_1$ . Therefore, there is a splitting

$$H_i(S^{2N+1} \times_{S^1} (LX/X)) = H_i(\mathbb{C}P^N) \oplus H_i((S^{2N+1} \times_{S^1} (LX/X))/p_2^{-1}(*))$$

generated by  $p_1$  and the collapsing map. Hence  $(p_2)_*$  is surjective.

Similarly any fixed point of the  $S^1$  action on  $LX$  defines a section of  $p'_1 : S^{2N+1} \times_{S^1} (LX) \rightarrow \mathbb{C}P^N$ . We consider the morphism of the Leray-Serre spectral sequences  $\pi_{*,*} : E_{*,*}(p'_1) \rightarrow E_{*,*}(p_1)$  of  $p_1$  and  $p'_1$  generated by  $\bar{q}$ .

By Proposition 2.4,  $H_i(LX/X; \mathbb{Q}) = 0$  for  $i < n - 2$  and  $n - 2 < i < 2n - 4$ . Note that since in the definition of  $X$  we collapse the  $(n - 2)$ -skeleton of  $B\pi$  instead of the  $(n - 1)$ -skeleton, the inequalities for  $i$  in Proposition 2.4 are shifted by one. Then the assumption  $n \geq 4$  implies that  $H_{n-1}(LX/X; \mathbb{Q}) = 0$  and hence,  $E_{2,n-1}^2(p_1) = 0$ . Therefore, nonzero elements on the  $(n + 1)$ -antidiagonal are only  $E_{0,n+1}^2$  and  $E_{n+1,0}^2$ . Since  $H_{n+1}(LX; \mathbb{Q}) \rightarrow H_{n+1}(LX/X; \mathbb{Q})$  is surjective,  $\phi_{0,n+1}^2 : E_{0,n}^2(p'_1) \rightarrow E_{0,n}^2(p_1)$  is surjective. Clearly,

$$\phi_{n+1,0}^2 : E_{n+1,0}^2(p'_1) = H_{n+1}(\mathbb{C}P^m; \mathbb{Q}) \rightarrow E_{n+1,0}^2(p_1) = H_{n+1}(\mathbb{C}P^m; \mathbb{Q})$$

is an isomorphism. By Proposition 2.5 part (a),  $\phi_{0,n+1}^\infty$  is surjective. Since both fibrations have sections, we have stabilizations:  $E_{n+1,0}^2(p_1) = E_{n+1,0}^\infty(p_1)$  and  $E_{n+1,0}^2(p'_1) = E_{n+1,0}^\infty(p'_1)$ . Then by Proposition 2.5 part (b),  $\bar{q}_*$  is surjective in dimension  $n + 1$ .  $\square$

### 3. THE MAIN RESULT

Let  $\nu : S^\infty \rightarrow \mathbb{C}P^\infty$  be the universal  $S^1$ -bundle. We identify

$$S^\infty = \{(E, v) \in \mathbb{C}P^\infty \times S^\infty \mid v \in E, |v| = 1\}$$

with an  $S^1$ -invariant subset of  $L(S^\infty)$  by sending  $(E, v)$  to the rotation  $f_v : S^1 \rightarrow E$  with  $f(1) = v$ . Then  $\mathbb{C}P^\infty$  is identified with a subset of  $S(S^\infty)$ . The same can be done for any principle  $S^1$ -bundle  $p : E \rightarrow B$ : The space  $E$  can be identified with an invariant subset of  $L(E)$  and  $B$  with a subset of  $S(E)$ . Let  $\hat{p} : B \rightarrow S(E)$  denote this embedding.

By  $cd(G)$  we denote the integral cohomological dimension of a group  $G$ . We recall that if  $cd(G) \neq 2$  then  $cd(G) = \min\{\dim K(G, 1)\}$  [Br].

**3.1. Theorem.** *Suppose that in the central extension*

$$1 \rightarrow C \rightarrow G \xrightarrow{p} H \rightarrow 1$$

where  $C \cong \mathbb{Z}$  the group  $G$  has finite  $cd(G) < \infty$  and  $H$  has property  $E$ . Then  $H_n(BH; \mathbb{Q}) = 0$  for  $n \geq \max\{cd(G) + 2, 5\}$ .

*Proof.* Assume that  $H_n(BH; \mathbb{Q}) \neq 0$  for some  $n \geq cd(G) + 2$ ,  $n \geq 5$ . Note that the homomorphism  $\phi : G \rightarrow H$  is the induced homomorphism  $p_* : \pi_1(BG) \rightarrow \pi_1(BH)$  for an oriented  $S^1$ -fibration  $p : BG \rightarrow BH$ . Since every oriented  $S^1$ -bundle is a principal  $S^1$ -bundle [Mo],  $p$  is the pull-back of the universal  $S^1$ -bundle  $\nu : S^\infty \rightarrow \mathbb{C}P^\infty$ . The composition  $S(p) \circ \hat{p} : BH \rightarrow S(BH)$  coincides with  $j_H$  where  $j_H : BH \rightarrow S_0(BH) \subset S(BH)$  is the inclusion map. Note that  $p : BG \rightarrow BH$  can be deformed to  $BH^{(n-2)}$  since  $BG$  is homotopy equivalent to a complex of dimension  $cd(G) \leq n - 2$ . Let  $H : BG \times I \rightarrow BH$  be a corresponding homotopy. It defines a homotopy  $\hat{H} : S(BG) \times I \rightarrow S(BH)$  and, hence a homotopy  $\tilde{H} : BH \times I \rightarrow S(BH)$  of  $S(p) \circ \hat{p}$ . Since  $\tilde{H}(BH \times \{0\}) \subset S_0(BH)$ , it follows that  $\tilde{H}(BH \times I) \subset S_0(BH)$ . Thus,  $\tilde{H}$  is a homotopy of  $j_H$  to a map  $g$  with image in  $S_0(BH^{(n-2)})$ . Hence we have a homotopy commutative diagram

$$\begin{array}{ccc} BH & \xleftarrow{id} & BH \\ g \downarrow & & j_H \downarrow \\ S_0(BH^{(n-2)}) & \xrightarrow{\subset} & S_0(BH) \end{array}$$

which defines a commutative diagram of homology groups

$$\begin{array}{ccc} H_n(BH; \mathbb{Q}) & \xleftarrow{id_*} & H_n(BH; \mathbb{Q}) \\ g_* \downarrow & & \cong \downarrow \\ H_n(S_0(BH^{(n-2)}); \mathbb{Q}) & \xrightarrow{0} & H_n(S_0(BH); \mathbb{Q}). \end{array}$$

The right vertical arrow is an isomorphism in view of Theorem 2.6 (a). By Theorem 2.6(b), the commutativity of the diagram implies that  $H_n(BH; \mathbb{Q}) = 0$ . This brings a contradiction.  $\square$

**Proof of Theorem 1.2.** The condition  $cd(G) < \infty$  implies that  $cd(G_x) < \infty$  for all  $x \in G$ . By Theorem 3.1 there is  $N \in \mathbb{N}$  such that  $H_m(N_x; \mathbb{Q}) = 0$  for  $m > N$  all  $x \in G$ . Therefore, the short exact sequence for  $T_*(x, \mathbb{Q})$ , (see the introduction) implies  $T_*(x, \mathbb{Q}) = 0$ .  $\square$

## REFERENCES

- [Ba] H. Bass, *Euler Characteristics and Characters of Discrete Groups*, Invent. math. 35 (1976), 155-196.
- [Bo] A. Borel, *Seminar on transformation groups*, Annals of mathematical Studies, vol 46, Princeton University Press, 1960.
- [Bre] G. Bredon, *Sheaf theory*, Second edition. Graduate Texts in Mathematics, 170. Springer-Verlag, New York, 1997.
- [Br] K. Brown, *Cohomology of groups*, Springer 1982.
- [Bu] D. Burghlea, *The cyclic homology of the group rings*, Comment. Math. Helvetici 60 (1985), 354-365.
- [CJY] R. Cohen, J. Jones, and J. Yan, *The loop homology algebra of spheres and projective spaces*, Categorical decomposition techniques in algebraic topology (Isle of Skye, 2001), 7792, Progr. Math., 215, Birkhuser, Basel, 2004.
- [C] Connes, Alain *Cohomologie cyclique et foncteurs Extn. (French) [Cyclic cohomology and functors Extn]* C. R. Acad. Sci. Paris Sr. I Math. 296 (1983), no. 23, 953-958.
- [E] B. Eckmann, *Cyclic homology of groups and the Bass conjecture*, Comment. Math. Helvetici 61 (1986), 193-202.
- [Em] I. Emmanouil, *On class of groups satisfying Bass' conjecture*, Invent. math. 132, (1998) 307-330.
- [EM] A. Engel, M. Marcinkowski, *Burghlea conjecture and asymptotic dimension of groups*. Preprint (2016) arXiv:1610.10076.
- [J] R. Ji, *Nilpotency of Connes' Periodicity Operator and the Idempotent Conjectures*, K-Theory 9 (1995), 59-76.
- [Li] P. A. Linnel, *Decomposition of augmentation ideals and relation modules*. Proc. London Math. Soc. (3) 47, (1983) 83-127.
- [L] Loday, Jean-Louis, *Cyclic homology*. Springer-Verlag, Berlin, 1992.
- [LQ] Loday, Jean-Louis; Quillen, Daniel *Homologie cyclique et homologie de l'algebre de Lie des matrices. (French) [Cyclic homology and homology of the Lie algebra of matrices]* C. R. Acad. Sci. Paris Sr. I Math. 296 (1983), no. 6, 295-297.
- [ML] S. MacLane, *Homology*. Reprint of the first edition. Die Grundlehren der mathematischen Wissenschaften, Band 114. Springer-Verlag, Berlin-New York, 1967.
- [Mo] Morita Shigeyuki, *Geometry of differential forms*. AMS 2001.
- [R] Rosenberg, Jonathan *Algebraic K-theory and its applications*. Graduate Texts in Mathematics, 147. Springer-Verlag, New York, 1994. x+392 pp.
- [T] Tsygan, B. L. *Homology of matrix Lie algebras over rings and the Hochschild homology*. (Russian) Uspekhi Mat. Nauk 38 (1983), no. 2(230), 217-218.
- [We] C. Westerland, *String homology of spheres and projective spaces* Algebr. Geom. Topol. 7 (2007), 309-325.

ALEXANDER N. DRANISHNIKOV, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF FLORIDA, 358 LITTLE HALL, GAINESVILLE, FL 32611-8105, USA  
*E-mail address:* dranish@math.ufl.edu